

Insights into electroreceptor development and evolution from molecular comparisons with hair cells

Clare V. H. Baker* and Melinda S. Modrell

Department of Physiology, Development & Neuroscience, University of Cambridge, Anatomy Building,
Downing Street, CB2 3DY, UK

From the symposium “Integrative Biology of Sensory Hair Cells” presented at the annual meeting of the
Society for Integrative and Comparative Biology, January 3–7, 2018 in San Francisco, California.

*Address for Correspondence:

Clare V. H. Baker

Department of Physiology, Development and Neuroscience

University of Cambridge

Anatomy Building, Downing Street, Cambridge CB2 3DY, United Kingdom

Tel +44 (0) 1223 333789

Fax +44 (0) 1223 333840

Email: cvhb1@cam.ac.uk

Running title: Electroreceptors and hair cells

Synopsis

The vertebrate lateral line system comprises a mechanosensory division, with neuromasts containing hair cells that detect local water movement ("distant touch"); and an electrosensory division, with electrosensory organs that detect the weak, low-frequency electric fields surrounding other animals in water (primarily used for hunting). The entire lateral line system was lost in the amniote lineage with the transition to fully terrestrial life; the electrosensory division was lost independently in several lineages, including the ancestors of frogs and of teleost fishes. (Electroreception with different characteristics subsequently evolved independently within two teleost lineages.) Recent gene expression studies in a non-teleost actinopterygian fish suggest that electroreceptor ribbon synapses employ the same transmission mechanisms as hair cell ribbon synapses, and show that developing electrosensory organs express transcription factors essential for hair cell development, including *Atoh1* and *Pou4f3*. Previous hypotheses for electroreceptor evolution suggest either that electroreceptors and hair cells evolved independently in the vertebrate ancestor from a common ciliated secondary cell, or that electroreceptors evolved from hair cells. The close developmental and putative physiological similarities implied by the gene expression data support the latter hypothesis, i.e., that electroreceptors evolved in the vertebrate ancestor as a "sister cell-type" to lateral line hair cells.

The mechanosensory division of the lateral line system

The sensory lateral line system (Coombs et al., 1989; Bullock et al., 2005; Coombs et al., 2014) only functions in water, and was lost in the lineage leading to amniotes; it is found in all fishes and aquatic-stage amphibians (except for myxiniid hagfishes and some direct-developing amphibians lacking an aquatic larval stage, in which it was lost secondarily; Bullock et al., 1983; Braun and Northcutt, 1997; Schlosser, 2002b). The mechanosensory division comprises lines of "neuromasts" along the flank (hence "lateral line") and around the head, often buried in canals that open to the water via pores (Webb, 2014). The sensory cells of neuromasts are hair cells, which morphologically resemble mammalian type 2 vestibular hair cells; like all hair cells, they are stimulated by deflection of the apical "hair bundle", i.e., a

staircase array of actin-rich microvilli ("stereocilia"), in the direction of the tallest stereocilia and adjacent primary cilium ("kinocilium") (Chagnaud and Coombs, 2014; Nicolson, 2017). This action triggers cation entry through a mechanically gated channel, whose identity is still controversial (Corey and Holt, 2016; Fettiplace, 2016; Wu and Müller, 2016) (see also Wu et al., 2017), depolarising the hair cell. This opens $\text{Ca}_v1.3$ channels (L-type voltage-gated Ca^{2+} channels) in the basolateral membrane, clustered beneath "pre-synaptic ribbons": proteinaceous structures that tether many glutamate-filled synaptic vesicles, enabling rapid, sustained neurotransmitter release at these "ribbon synapses" (Safieddine et al., 2012; Nicolson, 2015; Wichmann and Moser, 2015) with the nerve terminals of afferent lateral line neurons, whose cell bodies are collected in lateral line ganglia. Mechanosensory lateral line afferents project via the anterior or posterior lateral line nerves (depending on the position of the neuromasts) to the medial octavolateral nucleus in the hindbrain (Wullmann and Grothe, 2014). The mechanosensory lateral line responds to water movement relative to the body's surface, hence mediates a sense of "distant touch" (Dijkgraaf, 1963; Montgomery et al., 2014).

It has long been known from descriptive and fate-mapping studies in amphibian and, more recently, zebrafish embryos, that neuromasts and their afferent neurons develop from lateral line placodes - paired patches of thickened neurogenic ectoderm on the head (see e.g. Wright, 1951; Schlosser, 2002a; Piotrowski and Baker, 2014). Over 120 years ago, for example, Julia Platt memorably described watching the posterior lateral line placode of the urodele amphibian *Necturus* "plough its way through the skin, leaving a row of sense-organs in the wake" (Platt, 1896). Three developmentally and evolutionarily independent sets of paired cranial neurogenic placodes form hair cell-containing organs and their afferent neurons: the otic placode forms the inner ear (Alsina and Whitfield, 2017; Gálvez et al., 2017), the paired spiracular/paratympanic organ placode forms the anamniote spiracular organ and its amniote homologue (where present), the paratympanic organ (O'Neill et al., 2012), and the lateral line placodes form lateral line organs. All these placodes develop as thickenings within the "posterior placodal area" adjacent to the hindbrain (Grocott et al., 2012; Saint-Jeannet and Moody, 2014; Schlosser, 2014). Phylogenetic analysis suggests that the ancestor of all jawed vertebrates had three

lateral line placodes rostral to the otic placode, and three caudal to the otic placode (Northcutt, 1997). After forming neuroblasts, which delaminate and coalesce with neural crest-derived glia to form lateral line ganglia (which may in turn fuse with other nearby ganglia), lateral line placodes either elongate to form sensory ridges over the head, or migrate as a cell collective along the trunk, tracked by afferent axons and associated Schwann cells (Wright, 1951; Schlosser, 2002a; Piotrowski and Baker, 2014). The lateral line system of the teleost zebrafish is a widely used model for hair cell development and also regeneration, for example after treatment with ototoxic drugs such as aminoglycoside antibiotics (Thomas et al., 2015; Kniss et al., 2016; McGraw et al., 2017).

The electrosensory division of the lateral line system

Electroreception, i.e., the ability to detect weak electric fields in water, was only discovered in the mid-twentieth century, in conjunction with the finding that "weakly electric" teleost fishes generate electric fields (Lissmann, 1951; Lissmann and Machin, 1958; Lissmann, 1958) by discharging "electric organs" composed of modified muscle cells (Markham, 2013; Gallant et al., 2014). Weakly electric fishes have diverse "tuberous" electrosensory organs stimulated by the high-frequency electric fields generated by electric organ discharges, and morphologically distinct "ampullary" organs stimulated by the low-frequency electric fields around other animals (Bodznick and Montgomery, 2005; Jørgensen, 2005). (Tiny direct-current standing electric fields develop around animals in water owing to the leakage of ions across mucous membranes, with a low-frequency component imparted by, for example, rhythmic limb or ventilation movements; Wilkens and Hofmann, 2005; Bedore and Kajiura, 2013.) Teleost electroreceptors respond to anodal (exterior-positive) stimuli and are inhibited by cathodal stimuli, and are innervated (depending on their position) by either anterior or posterior lateral line nerves projecting to distinct "electrosensory lateral line lobes" in the medulla (Bullock et al., 1983; New, 1997; Alves-Gomes, 2001; Baker et al., 2013).

Following the discovery of electroreception in weakly electric teleosts, electroreception with different characteristics of physiology and innervation was discovered in representatives of all major

vertebrate groups: stimulation by weak, low-frequency cathodal (exterior-negative) electric fields, inhibition by weak anodal (exterior-positive) stimuli, and innervation by lateral line afferents projecting to the dorsal octavolateral nucleus in the medulla via the dorsal root of the anterior lateral line nerve (Bullock et al., 1983; New, 1997; Baker et al., 2013). Electoreception with all these characteristics is present in lampreys within the agnathan (jawless) fishes (which molecular evidence overwhelmingly supports as a monophyletic clade, the cyclostomes; Kuraku, 2008; Heimberg et al., 2010; Shimeld and Donoghue, 2012; Fig. 1), and across all three gnathostome (jawed) vertebrate groups: in all chondrichthyans (cartilaginous fishes), and in lineages within both groups of osteichthyans (bony fishes), i.e., in coelacanth, dipnoans (lungfishes) and both urodele and caecilian amphibians within the sarcopterygians (lobe-finned bony fishes and tetrapods), and in polypterids (bichirs, reedfishes) and chondrosteans (sturgeons, paddlefishes) within the actinopterygians (ray-finned bony fishes) (Bullock et al., 1983; New, 1997; Baker et al., 2013) (Fig. 1).

The most parsimonious hypothesis to explain this distribution is that electoreception was present in the ancestor of all living vertebrates, but lost independently in the agnathan lineage leading to hagfishes, in the sarcopterygian lineage leading to anuran amphibians, and in the actinopterygian lineage leading to the neopterygian fishes, i.e., holosteans (gars, bowfin) and teleosts (Bullock et al., 1983; New, 1997; Baker et al., 2013) (Fig. 1). Teleost electoreception is clearly different (see above), and only two related groups within each of two distinct lineages are electoreceptive, with one of the two groups in each lineage also having evolved electric organs and tuberous electoreceptors, likely representing multiple independent evolutionary events, probably via the diversification of neuromast hair cells (Bullock et al., 1983; New, 1997; Baker et al., 2013). (Within the amniotes, monotreme mammals and dolphins independently evolved trigeminal nerve-mediated electoreception, which will not be considered further here; for more information on this system, see Czech-Damal et al., 2013.)

Like hair cells, electoreceptors in all vertebrates have basolateral ribbon synapses with the terminals of their afferent neurons (Jørgensen, 2005). (Unlike hair cells, electoreceptors lack efferent innervation; Bodznick and Montgomery, 2005.) Electoreceptors in non-teleost fishes and amphibians

resemble immature hair cells (Northcutt et al., 1994) in having an apical primary cilium and variable numbers of microvilli (from none to hundreds, depending on the species) (Jørgensen, 2005), which are not organised into the stepped "hair bundle" that characterises mature hair cells (Lu and Sipe, 2016). In contrast, electroreceptors in the jawless lampreys and all teleosts bar one species have apical microvilli but no primary cilium (Jørgensen, 2005). (Lamprey electroreceptor development has not been characterised, however, so it is possible that a primary cilium forms but is lost during maturation, as in mammalian cochlear hair cells; Lu and Sipe, 2016.) (Nothing is known about the function of microvilli in electroreceptors, or why they should be present in some taxa but not others.) Lamprey electroreceptors are collected into superficial "end-bud" electrosensory organs, while in non-teleost gnathostomes, electroreceptors interspersed with supporting cells comprise the sensory epithelium in "ampullary organs": flask-shaped expansions at the base of a duct filled with conductive jelly (see e.g. Josberger et al., 2016), which opens to the surface via a pore (Jørgensen, 2005).

Over the last quarter-century, fate-mapping studies across the three major gnathostome groups have revealed that individual lateral line placodes form ampullary organs, as well as neuromasts and lateral line afferent neurons (Baker et al., 2013). These involved ablation and grafting studies in embryos of a sarcopterygian (a urodele amphibian, the axolotl *Ambystoma mexicanum*) (Northcutt et al., 1995), and vital dye (Dil)-labelling experiments in an actinopterygian (a chondrosteian, the Mississippi paddlefish *Polyodon spathula*) (Modrell et al., 2011) and a chondrichthyan (the little skate, *Leucoraja erinacea*) (Gillis et al., 2012). The placodes elongate to form sensory ridges, which ultimately fragment, with a line of neuromasts forming first, along the centre of the ridge, and ampullary organs forming later, from both flanks, such that fields of ampullary organs flank the neuromast line (Piotrowski and Baker, 2014). Furthermore, lateral line (and otic) placodes in all three species, as originally shown in another chondrichthyan, the shark *Scyliorhinus canicula*, express the transcriptional coactivator and phosphatase gene *Eya4*, and maintain *Eya4* expression in both electroreceptors and hair cells (O'Neill et al., 2007; Modrell et al., 2011; Gillis et al., 2012; Modrell and Baker, 2012).

Their shared embryonic origin in all three gnathostome groups, plus the shared characteristics of physiology and innervation as described earlier, together provide strong evidence for the homology of ampullary organs and electroreceptors across gnathostomes. Much less is known about agnathan electroreceptor development. Although sensory neurons derived from lamprey cranial placodes, including lateral line neurons, have been fate-mapped using focal Dil-labelling (Modrell et al., 2014), neither electroreceptors nor neuromasts have been identified at embryonic stages (though both are functional in ammocoete larvae; Ronan, 1988; Gelman et al., 2007). Thus, the embryonic origin of lamprey electroreceptors remains undetermined (as does that of teleost electroreceptors, though descriptive studies in a catfish suggest that ampullary organs form on the flanks of thin sensory ridges formed by elongating lateral line placodes, as in non-teleosts; see Northcutt, 2003; Baker et al., 2013). Nevertheless, the conserved physiology and innervation of lamprey and non-teleost gnathostome electroreceptors strongly supports their homology, i.e., that the electrosensory division of the lateral line system was inherited, like the mechanosensory division, from the common ancestor of all living vertebrates.

Molecular analysis suggests that electroreceptor synaptic transmission mechanisms are conserved with hair cells

As noted above, hair cells and electroreceptors are characterised by basolateral ribbon synapses with afferent nerve terminals, with presynaptic ribbons of varying morphologies (Jørgensen, 2005). The presynaptic ribbon is mainly composed of the only ribbon-specific protein known, Ribeye: this is produced from an alternative start site for the NAD(H)-sensitive transcriptional corepressor gene *Ctbp2*, resulting in an N-terminal A-domain found only in Ribeye (Schmitz et al., 2000; Stankiewicz et al., 2014). Vertebrate retinal and pineal photoreceptors, and retinal bipolar neurons, also have ribbon synapses (Zanazzi and Matthews, 2009; Matthews and Fuchs, 2010; Mercer and Thoreson, 2011; Safieddine et al., 2012). Presynaptic ribbons also contain conventional presynaptic cytomatrix proteins like Piccolo and Bassoon: in both photoreceptors and hair cells, the latter is found at the base of the ribbon, and is

required to anchor the ribbon to the plasma membrane (Zanazzi and Matthews, 2009; Matthews and Fuchs, 2010; Mercer and Thoreson, 2011; Safieddine et al., 2012). In the absence of Ribeye, the ribbon is lost and neurotransmitter release is reduced, but not abolished, at both retinal and hair cell synapses (Lv et al., 2016; Maxeiner et al., 2016; Becker et al., 2018; Jean et al., 2018). Thus, presynaptic ribbons may simply represent "an extreme example of the more general need for presynaptic machinery to facilitate the supply of release-ready vesicles to the active zone" (Matthews and Fuchs, 2010).

As described below, hair cell ribbon synapse physiology differs from other ribbon synapse physiology in multiple respects (Zanazzi and Matthews, 2009; Matthews and Fuchs, 2010; Mercer and Thoreson, 2011; Safieddine et al., 2012). Furthermore, as detailed below, our recent analysis of the expression of ribbon synapse-associated genes in a non-teleost actinopterygian suggests that electroreceptor ribbon synapses share the "unique" physiology of hair cell ribbon synapses (Modrell et al., 2017). These transcripts were identified in a set of just under 500 genes identified by differential RNA-seq as being enriched at least 1.85-fold in pooled operculum (gill-flap: covered in ampullary organs, plus some neuromasts) versus fin tissue (similar in tissue composition but without any lateral line organs) at late larval stages (the onset of independent feeding) in a chondrosteian, the paddlefish *Polyodon spathula* (Modrell et al., 2017).

In retinal photoreceptors, depolarisation elicits synaptic vesicle exocytosis by opening $\text{Ca}_v1.4$ channels (L-type voltage-gated Ca^{2+} channels) clustered near the presynaptic ribbon (Safieddine et al., 2012; Joiner and Lee, 2015). In hair cells, this role is played by $\text{Ca}_v1.3$ channels (whose pore-forming alpha subunit is encoded by *Cacna1d*) (Safieddine et al., 2012; Joiner and Lee, 2015; Nicolson, 2015), with the auxiliary subunit $\text{Ca}_v\beta2$ (encoded by *Cacnb2*) regulating their number and function (Neef et al., 2009). The only Ca^{2+} channel subunit genes present in the paddlefish lateral line-enriched gene-set were *Cacna1d* and *Cacnb2* (Modrell et al., 2017). *In situ* hybridisation (at the same late-larval stage used for RNA-seq) revealed that, in addition to expression in neuromasts, as expected (Nicolson, 2015; 2017), both *Cacna1d* and *Cacnb2* were expressed in ampullary organs (Fig. 2A-C; Modrell et al., 2017). This suggests that synaptic vesicle exocytosis is triggered in electroreceptors, as in hair cells, by $\text{Ca}_v1.3$

channels (using the same $\text{Ca}_v\beta 2$ subunit as in hair cells) rather than the $\text{Ca}_v1.4$ channels used at retinal photoreceptor ribbon synapses (Safieddine et al., 2012; Joiner and Lee, 2015). [Intriguingly, $\text{Ca}_v1.3$ was also recently identified as the apically located voltage-sensing Ca^{2+} channel responsible for electroreceptor function in a chondrichthyan, the little skate *Leucoraja erinacea* (Bellono et al., 2017), raising the possibility that $\text{Ca}_v1.3$ could also be the voltage-sensor for paddlefish electroreceptors (also see Modrell et al., 2017).]

In all ribbon synapses except hair cells, and at excitatory glutamatergic synapses in the central nervous system, the vesicular glutamate transporters Vglut1(encoded by *Slc17a7*) and/or Vglut2 (encoded by *Slc17a6*) are required to fill synaptic vesicles (Zanazzi and Matthews, 2009; Pangršič et al., 2012). In contrast, Vglut3 (encoded by *Slc17a8*) performs this function at hair cell ribbon synapses (Obholzer et al., 2008; Ruel et al., 2008; Seal et al., 2008). In the paddlefish lateral line-enriched gene-set, the only *Slc17a* gene family member present was *Slc17a8*, encoding Vglut3 (Modrell et al., 2017): this proved to be expressed in ampullary organs as well as neuromasts (Fig. 2D; Modrell et al., 2017). This finding suggests that in electroreceptors, as in hair cells but not in other ribbon synapse-containing cells, Vglut3 loads synaptic vesicles with neurotransmitter.

Furthermore, synaptic vesicle exocytosis is mediated at retinal ribbon synapses, as at other synapses, by neuronal SNAREs (soluble N-ethylmaleimide-sensitive fusion protein attachment protein receptors), whereas at hair cell ribbon synapses, SNAREs are replaced by a calcium-sensitive type II ferlin, otoferlin (Hams et al., 2017; Michalski et al., 2017) (also see Pangršič et al., 2012). The gene encoding otoferlin, *Otof*, was present in the paddlefish lateral line-enriched gene-set and expressed in ampullary organs as well as neuromasts (Fig. 2E; Modrell et al., 2017), suggesting that otoferlin mediates synaptic vesicle exocytosis at the ribbon synapses of both electroreceptors and hair cells.

Finally, although not unique to hair cell ribbon synapses, the paddlefish study identified the expression in ampullary organs (as well as neuromasts) of *Rims2*, encoding Rab3-interacting molecules 2 α and β , which are necessary to recruit $\text{Ca}_v1.3$ channels to the active zone membrane under the presynaptic ribbon of cochlear inner hair cells (Jung et al., 2015) (Fig. 2F; Modrell et al., 2017) and

Ribeye, identified by *in situ* hybridisation using a riboprobe against the N-terminal A-domain sequence of paddlefish *Ctbp2* (Fig. 2G; Modrell et al., 2017).

Taken together, these gene expression studies, albeit descriptive, suggest that the synaptic transmission mechanisms at the electroreceptor ribbon synapse are conserved with those of hair cells.

Developing ampullary organs express transcription factors essential for hair cell development, including *Atoh1* and *Pou4f3* (*Brn3c*)

The paddlefish lateral line-enriched gene-set included several transcription factor genes important for mammalian hair cell development (Modrell et al., 2017). Previous candidate gene approaches had already identified the expression of such genes in developing ampullary organs, as well as neuromasts, including the “pan-placodal” homeodomain transcription factor gene *Six1* and its transcription coactivator gene *Eya1* (Fig. 3A,B; Modrell et al., 2011). During cochlear hair cell development, *Six1* and *Eya1* interact physically with the high mobility group domain transcription factor *Sox2* to form a complex that is capable of inducing expression of the basic helix-loop-helix (bHLH) transcription factor gene *Atoh1* in cochlear explants (Ahmed et al., 2012; Zhang et al., 2017). *Atoh1* is necessary for hair cell formation (Bermingham et al., 1999) and is sufficient to promote hair cell differentiation in the context of the inner ear (Cai and Groves, 2015; Jahan et al., 2015; Costa et al., 2017). Both *Sox2* and *Atoh1* were also present in the paddlefish lateral line-enriched gene-set and proved to be expressed in developing ampullary organs, as well as neuromasts (Fig. 3C,D; Modrell et al., 2017). (*Atoh1* expression in developing ampullary organs was also reported briefly in a paper on paddlefish cerebellum development; Butts et al., 2014) Similarly, the POU domain transcription factor gene *Pou4f3* (*Brn3c*), which is essential for hair cell formation (Erkman et al., 1996; Xiang et al., 1997), and is induced in the cochlea by *Six1* and *Eya1*, independently of *Atoh1* (Ahmed et al., 2012), was in the paddlefish lateral line-enriched gene-set and expressed in developing ampullary organs (Fig. 3E; Modrell et al., 2017).

The expression of all these transcription factor genes in developing ampullary organs suggests that they are likely to be important for electroreceptor differentiation, not just hair cell differentiation.

Indeed, *Atoh1* is required for the differentiation of a variety of other cell types, including mechanosensory Merkel cells and proprioceptive neurons, plus cerebellar granule neurons and intestinal secretory cells: its precise role in directing hair cell development is surprisingly little understood (Cai and Groves, 2015; Jahan et al., 2015; Costa et al., 2017). Another bHLH transcription factor gene present in the paddlefish lateral line-enriched gene-set, *Neurod4*, may be important specifically for electroreceptor development, as its expression was restricted to developing ampullary organs, and never seen in neuromasts (Fig. 3F-H²; Modrell et al., 2017).

Although all these studies are descriptive, testing the function of these transcription factors and other genes during electroreceptor versus hair cell development is a very real prospect, given the efficiency of CRISPR/Cas9 genome-editing tools injected into fertilised eggs of urodele amphibians (so far, the axolotl and a newt, *Pleurodeles waltl*; see e.g. Flowers et al., 2014; Elewa et al., 2017). It may also be possible to target lateral line placodes directly as Cas9 protein/guide RNA complexes can also be electroporated into target tissue, resulting in very efficient genome editing (Fei et al., 2016). Such approaches could, in principle, be applied to any electroreceptive species for which embryos can be obtained frequently and in sufficiently large numbers to enable optimisation of CRISPR/Cas9 for that species.

Implications for electroreceptor evolution in the vertebrate ancestor

As proposed almost 40 years ago, based primarily on morphology (Jørgensen, 1982; 1989), electroreceptors could have evolved in the vertebrate ancestor via the diversification of already-evolved lateral line hair cells, or hair cells and electroreceptors could have been derived independently from a ciliated axon-less sensory cell, which itself probably evolved from an ancestral primary sensory neuron (see also Fritzsche and Straka, 2014). Indeed, putative hair cell homologues have been reported in tunicates (the closest invertebrate chordate relatives of the vertebrates; Fig. 1), as axon-less (secondary) sensory cells with one or more apical cilia and microvilli, afferent glutamatergic synapses, and conservation of some gene expression with hair cells (Burighel et al., 2003; 2011; Rigon et al., 2013;

2018), located in the coronal organ of the oral siphon tentacles and velum in ascidians, and in the circum-oral ring of appendicularians (Burighel et al., 2003; 2011; Rigon et al., 2013; 2018). Physiological evidence suggests that the ascidian coronal organ is mechanosensory (Mackie et al., 2006). It is plausible that the common ancestor of tunicates and vertebrates had scattered secondary mechanosensory cells, which evolved in the tunicate lineage into coronal organ cells in ascidians and circum-oral ring cells in appendicularians, and into hair cells (and electroreceptors) in vertebrates, even though the former (which are associated with oral regions) develop from more rostral ectoderm than vertebrate hair cells, which all develop from placodes forming from the posterior placodal area adjacent to the hindbrain (for a more detailed discussion, see Patthey et al., 2014; Schlosser et al., 2014; Schlosser, 2015).

However, the ribbon synapses that are characteristic of hair cells and electroreceptors (as well as retinal and pineal photoreceptors, and retinal bipolar neurons) seem to be vertebrate-specific: they have not been described in either tunicates (including the putative hair cell homologues; Burighel et al., 2003; Rigon et al., 2018) or the cephalochordates (amphioxus species) (see e.g. Petralia et al., 2015; 2016; 2017). Several independent lines of evidence support the homology of the frontal eye of amphioxus to the vertebrate retina (Lacalli, 1996; Vopalensky et al., 2012; Suzuki et al., 2015; Pergner and Kozmik, 2017), while the amphioxus lamellar body (located in the roof of the larval cerebral vesicle) is thought to be homologous to the vertebrate pineal organ (Eakin and Westfall, 1962; Ruiz and Anadón, 1991; Pergner and Kozmik, 2017). Thus, ribbon-less cellular homologues of retinal and, most likely, pineal photoreceptors were already present in separate locations within the central nervous system of the ancestral chordate. Taken together with the physiological differences between ribbon synapses in retinal versus hair cells (Zanazzi and Matthews, 2009; Matthews and Fuchs, 2010; Mercer and Thoreson, 2011; Safieddine et al., 2012), this suggests that ribbon synapse-containing cells could not all have evolved from a common ancestral cell type. Instead, the recruitment of Ribeye to the presynaptic active zone machinery, and thus the formation of presynaptic ribbons, must represent independent evolutionary events during the evolution of vertebrate retinal cells, pineal cells and hair cells. In all these sensory cell

types, increasing the number of synaptic vesicles available for ready release - the primary function of the ribbon (Lv et al., 2016; Maxeiner et al., 2016; Becker et al., 2018; Jean et al., 2018) - would have been a selective advantage.

The gene expression data from paddlefish suggest that ribbon synapses in electroreceptors use hair cell-specific transmission mechanisms (i.e., involving $Ca_v1.3$, $Vglut3$ and otoferlin; Fig. 2; Modrell et al., 2017). Similarly, the fact that so many transcription factor genes essential for hair cell differentiation, including *Atoh1* and *Pou4f3*, are also expressed in developing ampullary organs (Fig. 3; Modrell et al., 2011; 2017) suggests a very close developmental and evolutionary relationship with hair cells. Taking all these lines of evidence together, and with the caveat that nothing is yet known at the molecular level about electroreceptors in lampreys, it seems less likely that electroreceptors and hair cells evolved independently from a ciliated secondary sensory cell in the vertebrate ancestor, as this would also require electroreceptors to have recruited Ribeye to their presynaptic active zones, and formed presynaptic ribbons, independently of hair cells. Instead, given all the characteristics shared with hair cells, it seems more parsimonious to suggest that lateral line electroreceptors evolved in the vertebrate ancestor as a sister cell-type to lateral line hair cells (Arendt et al., 2016), i.e., via the diversification of lateral line hair cells.

Funding

This work was supported by the Biological and Biological Sciences Research Council (BB/F00818X/1 to C.V.H.B.), the Fisheries Society of the British Isles (Research Grant to M.S.M.) and the Leverhulme Trust (RPG-383 to C.V.H.B.).

References

Ahmed M, Wong EYM, Sun J, Xu J, Wang F, Xu P-X. 2012. Eya1-Six1 interaction is sufficient to induce hair cell fate in the cochlea by activating *Atoh1* expression in cooperation with Sox2. *Dev Cell* 22:377-390.

- Alsina B, Whitfield TT. 2017. Sculpting the labyrinth: Morphogenesis of the developing inner ear. *Semin Cell Dev Biol* 65:47-59.
- Alves-Gomes JA. 2001. The evolution of electroreception and bioelectrogenesis in teleost fish: a phylogenetic perspective. *J Fish Biol* 58:1489-1511.
- Arendt D, Musser JM, Baker CVH, Bergman A, Cepko C, Erwin DH, Pavlicev M, Schlosser G, Widder S, Laubichler MD, Wagner GP. 2016. The origin and evolution of cell types. *Nat Rev Genet* 17:744-757.
- Baker CVH, Modrell MS, Gillis JA. 2013. The evolution and development of vertebrate lateral line electroreceptors. *J Exp Biol* 216:2515-2522.
- Becker L, Schnee ME, Niwa M, Sun W, Maxeiner S, Talaei S, Kachar B, Rutherford MA, Ricci AJ. 2018. The presynaptic ribbon maintains vesicle populations at the hair cell afferent fiber synapse. *eLife* 7:e30241.
- Bedore CN, Kajiura SM. 2013. Bioelectric fields of marine organisms: voltage and frequency contributions to detectability by electroreceptive predators. *Physiol Biochem Zool* 86:298-311.
- Bellono NW, Leitch DB, Julius D. 2017. Molecular basis of ancestral vertebrate electroreception. *Nature* 543:391-396.
- Bermingham NA, Hassan BA, Price SD, Vollrath MA, Ben-Arie N, Eatock RA, Bellen HJ, Lysakowski A, Zoghbi HY. 1999. *Math1*: an essential gene for the generation of inner ear hair cells. *Science* 284:1837-1841.
- Bodznick D, Montgomery JC. 2005. The physiology of low-frequency electrosensory systems. In: Bullock TH, CD Hopkins, AN Popper, RR Fay, editors. *Electroreception*. New York: Springer. p. 132-53.
- Braun CB, Northcutt RG. 1997. The lateral line system of hagfishes (Craniata: Myxinoidea). *Acta Zool (Stockh)* 78:247-268.
- Bullock TH, Bodznick DA, Northcutt RG. 1983. The phylogenetic distribution of electroreception: evidence for convergent evolution of a primitive vertebrate sense modality. *Brain Res Rev* 287:25-46.

- Bullock TH, Hopkins CD, Popper AN, Fay RR, editors. 2005. Electoreception. New York: Springer.
- Burighel P, Lane NJ, Fabio G, Stefano T, Zaniolo G, Carnevali MD, Manni L. 2003. Novel, secondary sensory cell organ in ascidians: in search of the ancestor of the vertebrate lateral line. *J Comp Neurol* 461:236-249.
- Burighel P, Caicci F, Manni L. 2011. Hair cells in non-vertebrate models: lower chordates and molluscs. *Hear Res* 273:14-24.
- Butts T, Modrell MS, Baker CVH, Wingate RJT. 2014. The evolution of the vertebrate cerebellum: absence of a proliferative external granule layer in a non-teleost ray-finned fish. *Evol Dev* 16:92-100.
- Cai T, Groves AK. 2015. The role of Atonal factors in mechanosensory cell specification and function. *Mol Neurobiol* 52:1315-1329.
- Chagnaud BP, Coombs S. 2014. Information encoding and processing by the peripheral lateral line system. In: Coombs SC, H Bleckmann, RR Fay, AN Popper, editors. *The Lateral Line System*. New York: Springer. p. 151-94.
- Coombs S, Görner P, Münz H, editors. 1989. *The Mechanosensory Lateral Line. Neurobiology and Evolution*. New York: Springer-Verlag.
- Coombs SC, Bleckmann H, Fay RR, Popper AN, editors. 2014. *The Lateral Line System*. New York: Springer.
- Corey DP, Holt JR. 2016. Are TMCs the mechanotransduction channels of vertebrate hair cells? *J Neurosci* 36:10921-10926.
- Costa A, Powell LM, Lowell S, Jarman AP. 2017. Atoh1 in sensory hair cell development: constraints and cofactors. *Semin Cell Dev Biol* 65:60-68.
- Czech-Damal NU, Dehnhardt G, Manger P, Hanke W. 2013. Passive electroreception in aquatic mammals. *J Comp Physiol A* 199:555-563.
- Dijkgraaf S. 1963. The functioning and significance of the lateral-line organs. *Biol Rev* 38:51-105.
- Eakin RM, Westfall JA. 1962. Fine structure of photoreceptors in amphioxus. *J Ultrastruct Res* 6:531-539.

- Elewa A, Wang H, Talavera-López C, Joven A, Brito G, Kumar A, Hameed LS, Penrad-Mobayed M, Yao Z, Zamani N, Abbas Y, Abdullayev I, Sandberg R, Grabherr M, Andersson B, Simon A. 2017. Reading and editing the *Pleurodeles waltl* genome reveals novel features of tetrapod regeneration. Nat Commun 8:2286.
- Erkman L, McEvilly RJ, Luo L, Ryan AK, Hooshmand F, O'Connell SM, Keithley EM, Rapaport DH, Ryan AF, Rosenfeld MG. 1996. Role of transcription factors Brn-3.1 and Brn-3.2 in auditory and visual system development. Nature 381:603-606.
- Fei J-F, Knapp D, Schuez M, Murawala P, Zou Y, Singh SP, Drechsel D, Tanaka EM. 2016. Tissue-and time-directed electroporation of CAS9 protein–gRNA complexes *in vivo* yields efficient multigene knockout for studying gene function in regeneration. npj Regen Med 1:16002.
- Fettiplace R. 2016. Is TMC1 the hair cell mechanotransducer channel? Biophys J 111:3-9.
- Flowers GP, Timberlake AT, McLean KC, Monaghan JR, Crews CM. 2014. Highly efficient targeted mutagenesis in axolotl using Cas9 RNA-guided nuclease. Development 141:2165-2171.
- Fritzsche B, Straka H. 2014. Evolution of vertebrate mechanosensory hair cells and inner ears: toward identifying stimuli that select mutation driven altered morphologies. J Comp Physiol A 200:5-18.
- Gallant JR, Traeger LL, Volkening JD, Moffett H, Chen PH, Novina CD, Phillips GNJ, Anand R, Wells GB, Pinch M, Guth R, Unguez GA, Albert JS, Zakon HH, Samanta MP, Sussman MR. 2014. Genomic basis for the convergent evolution of electric organs. Science 344:1522-1525.
- Gálvez H, Abelló G, Giraldez F. 2017. Signaling and transcription factors during inner ear development: the generation of hair cells and otic neurons. Front Cell Dev Biol 5:21.
- Gelman S, Ayali A, Tytell ED, Cohen AH. 2007. Larval lampreys possess a functional lateral line system. J Comp Physiol A 193:271-277.
- Gillis JA, Modrell MS, Northcutt RG, Catania KC, Luer CA, Baker CVH. 2012. Electrosensory ampullary organs are derived from lateral line placodes in cartilaginous fishes. Development 139:3142-3146.
- Grocott T, Tambalo M, Streit A. 2012. The peripheral sensory nervous system in the vertebrate head: a gene regulatory perspective. Dev Biol 370:3-23.

- Hams N, Padmanarayana M, Qiu W, Johnson CP. 2017. Otoferlin is a multivalent calcium-sensitive scaffold linking SNAREs and calcium channels. *Proc Natl Acad Sci USA* 114:8023-8028.
- Heimberg AM, Cowper-Sal Lari R, Semon M, Donoghue PC, Peterson KJ. 2010. MicroRNAs reveal the interrelationships of hagfish, lampreys, and gnathostomes and the nature of the ancestral vertebrate. *Proc Natl Acad Sci USA* 107:19379-19383.
- Jahan I, Pan N, Fritzsche B. 2015. Opportunities and limits of the one gene approach: the ability of Atoh1 to differentiate and maintain hair cells depends on the molecular context. *Front Cell Neurosci* 9:26.
- Jean P, Lopez de la Morena D, Michanski S, Jaime Tobón LM, Chakrabarti R, Picher MM, Neef J, Jung S, Gültas M, Maxeiner S, Neef A, Wichmann C, Strenzke N, Grabner C, Moser T. 2018. The synaptic ribbon is critical for sound encoding at high rates and with temporal precision. *eLife* 7:e29275.
- Joiner ML, Lee A. 2015. Voltage-gated Ca_v1 channels in disorders of vision and hearing. *Curr Mol Pharmacol* 8:143-148.
- Jørgensen JM. 1982. Fine structure of the ampullary organs of the bichir *Polypterus senegalus* Cuvier, 1829 (Pisces: Brachiopterygii) with some notes on the phylogenetic development of electroreceptors. *Acta Zool (Stockh)* 63:211-217.
- Jørgensen JM. 1989. Evolution of octavolateralis sensory cells. In: Coombs S, P Görner, H Münz, editors. *The Mechanosensory Lateral Line. Neurobiology and Evolution*. New York: Springer-Verlag. p. 115-45.
- Jørgensen JM. 2005. Morphology of electroreceptive sensory organs. In: Bullock TH, CD Hopkins, AN Popper, RR Fay, editors. *Electroreception*. New York: Springer. p. 47-67.
- Josberger EE, Hassanzadeh P, Deng Y, Sohn J, Rego MJ, Amemiya CT, Rolandi M. 2016. Proton conductivity in ampullae of Lorenzini jelly. *Sci Adv* 2:e1600112.
- Jung S, Oshima-Takago T, Chakrabarti R, Wong AB, Jing Z, Yamanbaeva G, Picher MM, Wojcik SM, Göttfert F, Predoehl F, Michel K, Hell SW, Schoch S, Strenzke N, Wichmann C, Moser T. 2015. Rab3-interacting molecules 2 α and 2 β promote the abundance of voltage-gated Ca_v1.3 Ca²⁺

- channels at hair cell active zones. *Proc Natl Acad Sci USA* 112:E3141-9.
- Kniss JS, Jiang L, Piotrowski T. 2016. Insights into sensory hair cell regeneration from the zebrafish lateral line. *Curr Opin Genet Dev* 40:32-40.
- Kuraku S. 2008. Insights into cyclostome phylogenomics: pre-2R or post-2R? *Zoolog Sci* 25:960-968.
- Lacalli TC. 1996. Frontal eye circuitry, rostral sensory pathways and brain organization in amphioxus larvae: evidence from 3D reconstructions. *Phil Trans R Soc Lond B* 351:243-263.
- Lissmann HW. 1951. Continuous electrical signals from the tail of a fish, *Gymnarchus niloticus* Cuv. *Nature* 167:201-202.
- Lissmann HW. 1958. On the function and evolution of electric organs in fish. *J Exp Biol* 35:156-191.
- Lissmann HW, Machin KE. 1958. The mechanism of object location in *Gymnarchus niloticus* and similar fish. *J Exp Biol* 35:451-486.
- Lu X, Sipe CW. 2016. Developmental regulation of planar cell polarity and hair-bundle morphogenesis in auditory hair cells: lessons from human and mouse genetics. *WIREs Dev Biol* 5:85-101.
- Lv C, Stewart WJ, Akanyeti O, Frederick C, Zhu J, Santos-Sacchi J, Sheets L, Liao JC, Zenisek D. 2016. Synaptic ribbons require Ribeye for electron density, proper synaptic localization, and recruitment of calcium channels. *Cell Rep* 15:2784-2795.
- Mackie GO, Burighel P, Caicci F, Manni L. 2006. Innervation of ascidian siphons and their responses to stimulation. *Can J Zool* 84:1146-1162.
- Markham MR. 2013. Electrocyte physiology: 50 years later. *J Exp Biol* 216:2451-2458.
- Matthews G, Fuchs P. 2010. The diverse roles of ribbon synapses in sensory neurotransmission. *Nat Rev Neurosci* 11:812-822.
- Maxeiner S, Luo F, Tan A, Schmitz F, Südhof TC. 2016. How to make a synaptic ribbon: RIBEYE deletion abolishes ribbons in retinal synapses and disrupts neurotransmitter release. *EMBO J* 35:1098-1114.
- McGraw HF, Drerup CM, Nicolson T, Nechiporuk AV. 2017. The molecular and cellular mechanisms of zebrafish lateral line development. In: Cramer KS, AB Coffin, RR Fay, AN Popper, editors. *Auditory*

Development and Plasticity. Springer International Publishing p. 49-73.

- Mercer AJ, Thoreson WB. 2011. The dynamic architecture of photoreceptor ribbon synapses: cytoskeletal, extracellular matrix, and intramembrane proteins. *Vis Neurosci* 28:453-471.
- Michalski N, Goutman JD, Auclair SM, Boutet de Monvel J, Tertrais M, Emptoz A, Parrin A, Nouaille S, Guillon M, Sachse M, Ciric D, Bahloul A, Hardelin JP, Sutton RB, Avan P, Krishnakumar SS, Rothman JE, Dulon D, Safieddine S, Petit C. 2017. Otoferlin acts as a Ca^{2+} sensor for vesicle fusion and vesicle pool replenishment at auditory hair cell ribbon synapses. *eLife* 6:e31013.
- Modrell MS, Baker CVH. 2012. Evolution of electrosensory ampullary organs: conservation of *Eya4* expression during lateral line development in jawed vertebrates. *Evol Dev* 14:277-285.
- Modrell MS, Bemis WE, Northcutt RG, Davis MC, Baker CVH. 2011. Electrosensory ampullary organs are derived from lateral line placodes in bony fishes. *Nat Commun* 2:496.
- Modrell MS, Hockman D, Uy B, Buckley D, Sauka-Spengler T, Bronner ME, Baker CVH. 2014. A fate-map for cranial sensory ganglia in the sea lamprey. *Dev Biol* 385:405-416.
- Modrell MS, Lyne M, Carr AR, Zakon HH, Buckley D, Campbell AS, Davis MC, Micklem G, Baker CVH. 2017. Insights into electrosensory organ development, physiology and evolution from a lateral line-enriched transcriptome. *eLife* 6:e24197.
- Montgomery J, Bleckmann H, Coombs S. 2014. Sensory ecology and neuroethology of the lateral line. In: Coombs SC, H Bleckmann, RR Fay, AN Popper, editors. *The Lateral Line System*. New York: Springer. p. 121-50.
- Neef J, Gehrt A, Bulankina AV, Meyer AC, Riedel D, Gregg RG, Strenzke N, Moser T. 2009. The Ca^{2+} channel subunit $\beta 2$ regulates Ca^{2+} channel abundance and function in inner hair cells and is required for hearing. *J Neurosci* 29:10730-10740.
- New JG. 1997. The evolution of vertebrate electrosensory systems. *Brain Behav Evol* 50:244-252.
- Nicolson T. 2015. Ribbon synapses in zebrafish hair cells. *Hear Res* 330:170-177.
- Nicolson T. 2017. The genetics of hair-cell function in zebrafish. *J Neurogenet* 31:102-112.
- Northcutt RG. 1997. Evolution of gnathostome lateral line ontogenies. *Brain Behav Evol* 50:25-37.

- Northcutt RG. 2003. Development of the lateral line system in the channel catfish. In: Browman HI AB Skiftesvik, editors. The Big Fish Bang. Proceedings of the 26th Annual Larval Fish Conference. Bergen, Norway: Institute of Marine Research. p. 137-59.
- Northcutt RG, Catania KC, Criley BB. 1994. Development of lateral line organs in the axolotl. *J Comp Neurol* 340:480-514.
- Northcutt RG, Brändle K, Fritsch B. 1995. Electoreceptors and mechanosensory lateral line organs arise from single placodes in axolotls. *Dev Biol* 168:358-373.
- O'Neill P, McCole RB, Baker CVH. 2007. A molecular analysis of neurogenic placode and cranial sensory ganglion development in the shark, *Scyliorhinus canicula*. *Dev Biol* 304:156-181.
- O'Neill P, Mak S-S, Fritsch B, Ladher RK, Baker CVH. 2012. The amniote paratympanic organ develops from a previously undiscovered sensory placode. *Nat Commun* 3:1041.
- Obholzer N, Wolfson S, Trapani JG, Mo W, Nechiporuk A, Busch-Nentwich E, Seiler C, Sidi S, Söllner C, Duncan RN, Boehland A, Nicolson T. 2008. Vesicular glutamate transporter 3 is required for synaptic transmission in zebrafish hair cells. *J Neurosci* 28:2110-2118.
- Pangršič T, Reisinger E, Moser T. 2012. Otoferlin: a multi-C2 domain protein essential for hearing. *Trends Neurosci* 35:671-680.
- Patthey C, Schlosser G, Shimeld SM. 2014. The evolutionary history of vertebrate cranial placodes - I: cell type evolution. *Dev Biol* 389:82-97.
- Pergner J, Kozmik Z. 2017. Amphioxus photoreceptors - insights into the evolution of vertebrate opsins, vision and circadian rhythmicity. *Int J Dev Biol* 61:665-681.
- Petralia RS, Wang Y-X, Mattson MP, Yao PJ. 2015. Structure, distribution, and function of neuronal/synaptic spinules and related invaginating projections. *NeuroMolecular Med* 17:211-240.
- Petralia RS, Wang Y-X, Mattson MP, Yao PJ. 2016. The diversity of spine synapses in animals. *NeuroMolecular Med* 18:497-539.
- Petralia RS, Wang Y-X, Mattson MP, Yao PJ. 2017. Invaginating presynaptic terminals in neuromuscular junctions, photoreceptor terminals, and other synapses of animals. *NeuroMolecular Med* 19:193-

- Piotrowski T, Baker CVH. 2014. The development of lateral line placodes: Taking a broader view. *Dev Biol* 389:68-81.
- Platt JB. 1896. Ontogenetic differentiation of the ectoderm in *Necturus*. Study II. On the development of the peripheral nervous system. *Quart J Microsc Sci* 38:485-547.
- Rigon F, Stach T, Caicci F, Gasparini F, Burighel P, Manni L. 2013. Evolutionary diversification of secondary mechanoreceptor cells in tunicata. *BMC Evol Biol* 13:112.
- Rigon F, Gasparini F, Shimeld SM, Candiani S, Manni L. 2018. Developmental signature, synaptic connectivity and neurotransmission are conserved between vertebrate hair cells and tunicate coronal cells. *J Comp Neurol* 526:957-971.
- Ronan M. 1988. Anatomical and physiological evidence for electroreception in larval lampreys. *Brain Res* 448:173-177.
- Ruel J, Emery S, Nouvian R, Bersot T, Amilhon B, Van Rybroek JM, Rebillard G, Lenoir M, Eybalin M, Delprat B, Sivakumaran TA, Giros B, El Mestikawy S, Moser T, Smith RJH, Lesperance MM, Puel J-L. 2008. Impairment of *SLC17A8* encoding vesicular glutamate transporter-3, VGLUT3, underlies nonsyndromic deafness DFNA25 and inner hair cell dysfunction in null mice. *Am J Hum Genet* 83:278-292.
- Ruiz S, Anadón R. 1991. The fine structure of lamellate cells in the brain of amphioxus (*Branchiostoma lanceolatum*, Cephalochordata). *Cell Tissue Res* 263:597-600.
- Safieddine S, El-Amraoui A, Petit C. 2012. The auditory hair cell ribbon synapse: from assembly to function. *Annu Rev Neurosci* 35:509-528.
- Saint-Jeannet J-P, Moody SA. 2014. Establishing the pre-placodal region and breaking it into placodes with distinct identities. *Dev Biol* 389:13-27.
- Schlosser G. 2002a. Development and evolution of lateral line placodes in amphibians I. Development. *Zoology (Jena)* 105:119-146.
- Schlosser G. 2002b. Development and evolution of lateral line placodes in amphibians. II. Evolutionary

- diversification. *Zoology (Jena)* 105:177-193.
- Schlosser G. 2014. Early embryonic specification of vertebrate cranial placodes. *WIREs Dev Biol* 3:349-363.
- Schlosser G. 2015. Vertebrate cranial placodes as evolutionary innovations - the ancestor's tale. *Curr Top Dev Biol* 111:235-300.
- Schlosser G, Patthey C, Shimeld SM. 2014. The evolutionary history of vertebrate cranial placodes II. Evolution of ectodermal patterning. *Dev Biol* 389:98-119.
- Schmitz F, Königstorfer A, Südhof TC. 2000. RIBEYE, a component of synaptic ribbons: a protein's journey through evolution provides insight into synaptic ribbon function. *Neuron* 28:857-872.
- Seal RP, Akil O, Yi E, Weber CM, Grant L, Yoo J, Clause A, Kandler K, Noebels JL, Glowatzki E, Lustig LR, Edwards RH. 2008. Sensorineural deafness and seizures in mice lacking vesicular glutamate transporter 3. *Neuron* 57:263-275.
- Shimeld SM, Donoghue PCJ. 2012. Evolutionary crossroads in developmental biology: cyclostomes (lamprey and hagfish). *Development* 139:2091-2099.
- Stankiewicz TR, Gray JJ, Winter AN, Linseman DA. 2014. C-terminal binding proteins: central players in development and disease. *Biomol Concepts* 5:489-511.
- Suzuki DG, Murakami Y, Escriva H, Wada H. 2015. A comparative examination of neural circuit and brain patterning between the lamprey and amphioxus reveals the evolutionary origin of the vertebrate visual center. *J Comp Neurol* 523:251-261.
- Thomas ED, Cruz IA, Hailey DW, Raible DW. 2015. There and back again: development and regeneration of the zebrafish lateral line system. *WIREs Dev Biol* 4:1-16.
- Vopalensky P, Pergner J, Liegertova M, Benito-Gutierrez E, Arendt D, Kozmik Z. 2012. Molecular analysis of the amphioxus frontal eye unravels the evolutionary origin of the retina and pigment cells of the vertebrate eye. *Proc Natl Acad Sci USA* 109:15383-15388.
- Webb JF. 2014. Morphological diversity, development, and evolution of the mechanosensory lateral line system. In: Coombs SC, H Bleckmann, RR Fay, AN Popper, editors. *The Lateral Line System*. New

York: Springer. p. 17-72.

Wichmann C, Moser T. 2015. Relating structure and function of inner hair cell ribbon synapses. *Cell Tissue Res* 361:95-114.

Wilkens LA, Hofmann MH. 2005. Behaviour of animals with passive, low-frequency electrosensory systems. In: Bullock TH, CD Hopkins, AN Popper, RR Fay, editors. *Electroreception*. New York: Springer. p. 229-63.

Wright MR. 1951. The lateral line system of sense organs. *Q Rev Biol* 26:264-280.

Wu Z, Müller U. 2016. Molecular identity of the mechanotransduction channel in hair cells: not quiet there yet. *J Neurosci* 36:10927-10934.

Wu Z, Grillet N, Zhao B, Cunningham C, Harkins-Perry S, Coste B, Ranade S, Zebarjadi N, Beurg M, Fettiplace R, Patapoutian A, Müller U. 2017. Mechanosensory hair cells express two molecularly distinct mechanotransduction channels. *Nat Neurosci* 20:24-33.

Wullimann MF, Grothe B. 2014. The central nervous organization of the lateral line system. In: Coombs SC, H Bleckmann, RR Fay, AN Popper, editors. *The Lateral Line System*. New York: Springer. p. 195-251.

Xiang M, Gan L, Li D, Chen Z-Y, Zhou L, O'Malley BW, Jr., Klein W, Nathans J. 1997. Essential role of POU-domain factor Brn-3c in auditory and vestibular hair cell development. *Proc Natl Acad Sci USA* 94:9445-9450.

Zanazzi G, Matthews G. 2009. The molecular architecture of ribbon presynaptic terminals. *Mol Neurobiol* 39:130-148.

Zhang T, Xu J, Maire P, Xu P-X. 2017. Six1 is essential for differentiation and patterning of the mammalian auditory sensory epithelium. *PLoS Genet* 13:e1006967.

Figure Legends

Fig. 1 Phylogeny showing the distribution of lateral line sensory divisions among living vertebrates, with the invertebrate chordates shown for reference. Black font indicates possession of both the mechanosensory division and the electrosensory division (the latter responding to low-frequency, cathodal stimuli, with the dorsal octavolateral nucleus as the hindbrain target of lateral line afferents projecting via the dorsal root of the anterior lateral line nerve). Grey font indicates possession of the mechanosensory division only, except for the amniotes (☼), which lost the whole lateral line system during the transition to fully terrestrial life, and actinopterygian teleost fishes (≠), in which a few groups independently evolved lateral line electroreception responding to anodal stimuli, with an electrosensory lateral line lobe as the hindbrain target of lateral line afferents projecting via both anterior and posterior lateral line nerves.

Fig. 2 Genes important for transmission at the hair cell ribbon synapse are expressed in electrosensory organs in the paddlefish. (A) Schematic showing the distribution of neuromasts in canals, flanked by large fields of electrosensory ampullary organs, on the head of a late-larval paddlefish (at the onset of independent feeding). The caudal box outlines the region shown at higher power in panels C,D; the rostral box outlines the region shown at higher power in panels E-G. (B-G) *In situ* hybridisation (shown in skin-mounts in panels C-G: white dotted lines indicating the approximate position of lateral line canals, within which neuromasts are buried) reveals expression of hair cell ribbon synapse-associated genes in ampullary organs, as well as neuromasts, for (B) *Cacna1d*, encoding the pore-forming subunit of Ca_v1.3; (C) *Cacnb2*, encoding the Ca_vβ2 auxiliary subunit for Ca_v1.3 in hair cells (expression is very weak in neuromasts: compare with panel D for neuromast distribution); (D) *Sc17a8*, encoding Vglut3 (asterisk indicates damage to the skin-mount); (E) *Otof*, encoding otoferlin; (F) *Rims2*, encoding Rab3-interacting molecules 2 α and β, which recruit Ca_v1.3 channels to the membrane under the presynaptic ribbon in hair cells; (G) the N-terminal A-domain sequence of *Ctbp2*, encoding Ribeye. Scale-bars: 200 μm in B; 100 μm in C (panels D-G are shown at the same magnification as panel C). Abbreviations: ao, ampullary organs; e, eye; io, infraorbital neuromast line; nm, neuromasts; ol, otic neuromast line; so, supraorbital neuromast line. (A colour version of this figure is available online.)

Fig. 3 Transcription factor genes important for hair cell development are expressed in developing electrosensory organs in the paddlefish. (A-E) *In situ* hybridisation in paddlefish embryos reveals expression in both developing neuromasts and ampullary organs of (A) *Six1* and (B) *Eya1* (expression of both is also seen in the posterior lateral line primordium migrating along the trunk); (C) *Sox2*, (D) *Atoh1* (the inset shows neuromast lines at an earlier stage of development) and (E) *Pou4f3* (*Bmn3c*). **(F-H²)** The bHLH transcription factor gene *Neurod4* is expressed in ampullary organs but not neuromasts. The dotted white lines in panel G show the approximate position of the neuromast lines. H-H² show a transverse section through the head (near the eye). Panel H shows *Neurod4* expression in an ampullary organ. Panel H¹ shows immunostaining for a calcium-buffering protein (an oncomodulin-related beta-parvalbumin; see Modrell et al., 2017) expressed in both neuromast hair cells and electroreceptors, which shows a *Neurod4*-negative neuromast dorsal to a *Neurod4*-positive ampullary organ. Panel H² shows DAPI, which stains all nuclei. Scale-bars: 500 μ m in A (panels B-E shown at the same magnification as panel A); 500 μ m in F; 100 μ m in G; 50 μ m in H. Abbreviations: ao, ampullary organ; e, eye; nm, neuromasts; pll, posterior lateral line primordium. (A colour version of this figure is available online.)





